

FIGURE 1: Recommended immunization schedule for persons aged 0 through 6 years—United States, 2012 (for those who fall behind or start late, see the catch-up schedule [Figure 3])

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	9 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years	
Hepatitis B ¹	Hep B	HepB												Range of recommended ages for all children
Rotavirus ²				RV	RV	RV ²								
Diphtheria, tetanus, pertussis ³				DTaP	DTaP	DTaP	see footnote ³		DTaP				DTaP	
<i>Haemophilus influenzae</i> type b ⁴				Hib	Hib	Hib ⁴		Hib						Range of recommended ages for certain high-risk groups
Pneumococcal ⁵				PCV	PCV	PCV		PCV				PPSV		
Inactivated poliovirus ⁶				IPV	IPV			IPV					IPV	
Influenza ⁷														
Measles, mumps, rubella ⁸								MMR		see footnote ⁸			MMR	Range of recommended ages for all children and certain high-risk groups
Varicella ⁹								Varicella		see footnote ⁹			Varicella	
Hepatitis A ¹⁰														
Meningococcal ¹¹														

This schedule includes recommendations in effect as of December 23, 2011. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Vaccination providers should consult the relevant Advisory Committee on Immunization Practices (ACIP) statement for detailed recommendations, available online at <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow vaccination should be reported to the Vaccine Adverse Event Reporting System (VAERS) online (<http://www.vaers.hhs.gov>) or by telephone (800-822-7967).

- Hepatitis B (HepB) vaccine.** (Minimum age: birth)
At birth:
 - Administer monovalent HepB vaccine to all newborns before hospital discharge.
 - For infants born to hepatitis B surface antigen (HBsAg)-positive mothers, administer HepB vaccine and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth. These infants should be tested for HBsAg and antibody to HBsAg (anti-HBs) 1 to 2 months after receiving the last dose of the series.
 - If mother's HBsAg status is unknown, within 12 hours of birth administer HepB vaccine for infants weighing $\geq 2,000$ grams, and HepB vaccine plus HBIG for infants weighing $< 2,000$ grams. Determine mother's HBsAg status as soon as possible and, if she is HBsAg-positive, administer HBIG for infants weighing $\geq 2,000$ grams (no later than age 1 week).**Doses after the birth dose:**
 - The second dose should be administered at age 1 to 2 months. Monovalent HepB vaccine should be used for doses administered before age 6 weeks.
 - Administration of a total of 4 doses of HepB vaccine is permissible when a combination vaccine containing HepB is administered after the birth dose.
 - Infants who did not receive a birth dose should receive 3 doses of a HepB-containing vaccine starting as soon as feasible (Figure 3).
 - The minimum interval between dose 1 and dose 2 is 4 weeks, and between dose 2 and 3 is 8 weeks. The final (third or fourth) dose in the HepB vaccine series should be administered no earlier than age 24 weeks and at least 16 weeks after the first dose.
- Rotavirus (RV) vaccines.** (Minimum age: 6 weeks for both RV-1 [Rotarix] and RV-5 [Rota Teq])
 - The maximum age for the first dose in the series is 14 weeks, 6 days; and 8 months, 0 days for the final dose in the series. Vaccination should not be initiated for infants aged 15 weeks, 0 days or older.
 - If RV-1 (Rotarix) is administered at ages 2 and 4 months, a dose at 6 months is not indicated.
- Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine.** (Minimum age: 6 weeks)
 - The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.
- Haemophilus influenzae* type b (Hib) conjugate vaccine.** (Minimum age: 6 weeks)
 - If PRP-OMP (PedvaxHIB or Comvax [HepB-Hib]) is administered at ages 2 and 4 months, a dose at age 6 months is not indicated.
 - Hiberix should only be used for the booster (final) dose in children aged 12 months through 4 years.
- Pneumococcal vaccines.** (Minimum age: 6 weeks for pneumococcal conjugate vaccine [PCV]; 2 years for pneumococcal polysaccharide vaccine [PPSV])
 - Administer 1 dose of PCV to all healthy children aged 24 through 59 months who are not completely vaccinated for their age.
 - For children who have received an age-appropriate series of 7-valent PCV (PCV7), a single supplemental dose of 13-valent PCV (PCV13) is recommended for:
 - All children aged 14 through 59 months
 - Children aged 60 through 71 months with underlying medical conditions.
 - Administer PPSV at least 8 weeks after last dose of PCV to children aged 2 years or older with certain underlying medical conditions, including a cochlear implant. See *MMWR* 2010;59(No. RR-11), available at <http://www.cdc.gov/mmwr/pdf/rr/rr5911.pdf>.
- Inactivated poliovirus vaccine (IPV).** (Minimum age: 6 weeks)
 - If 4 or more doses are administered before age 4 years, an additional dose should be administered at age 4 through 6 years.
 - The final dose in the series should be administered on or after the fourth birthday and at least 6 months after the previous dose.
- Influenza vaccines.** (Minimum age: 6 months for trivalent inactivated influenza vaccine [TIV]; 2 years for live, attenuated influenza vaccine [LAIV])
 - For most healthy children aged 2 years and older, either LAIV or TIV may be used. However, LAIV should not be administered to some children, including 1) children with asthma, 2) children 2 through 4 years who had wheezing in the past 12 months, or 3) children who have any other underlying medical conditions that predispose them to influenza complications. For all other contraindications to use of LAIV, see *MMWR* 2010;59(No. RR-8), available at <http://www.cdc.gov/mmwr/pdf/rr/rr5908.pdf>.
 - For children aged 6 months through 8 years:
 - For the 2011–12 season, administer 2 doses (separated by at least 4 weeks) to those who did not receive at least 1 dose of the 2010–11 vaccine. Those who received at least 1 dose of the 2010–11 vaccine require 1 dose for the 2011–12 season.
 - For the 2012–13 season, follow dosing guidelines in the 2012 ACIP influenza vaccine recommendations.
- Measles, mumps, and rubella (MMR) vaccine.** (Minimum age: 12 months)
 - The second dose may be administered before age 4 years, provided at least 4 weeks have elapsed since the first dose.
 - Administer MMR vaccine to infants aged 6 through 11 months who are traveling internationally. These children should be revaccinated with 2 doses of MMR vaccine, the first at ages 12 through 15 months and at least 4 weeks after the previous dose, and the second at ages 4 through 6 years.
- Varicella (VAR) vaccine.** (Minimum age: 12 months)
 - The second dose may be administered before age 4 years, provided at least 3 months have elapsed since the first dose.
 - For children aged 12 months through 12 years, the recommended minimum interval between doses is 3 months. However, if the second dose was administered at least 4 weeks after the first dose, it can be accepted as valid.
- Hepatitis A (HepA) vaccine.** (Minimum age: 12 months)
 - Administer the second (final) dose 6 to 18 months after the first.
 - Unvaccinated children 24 months and older at high risk should be vaccinated. See *MMWR* 2006;55(No. RR-7), available at <http://www.cdc.gov/mmwr/pdf/rr/rr5507.pdf>.
 - A 2-dose HepA vaccine series is recommended for anyone aged 24 months and older, previously unvaccinated, for whom immunity against hepatitis A virus infection is desired.
- Meningococcal conjugate vaccines, quadrivalent (MCV4).** (Minimum age: 9 months for Menactra [MCV4-D], 2 years for Menveo [MCV4-CRM])
 - For children aged 9 through 23 months 1) with persistent complement component deficiency; 2) who are residents of or travelers to countries with hyperendemic or epidemic disease; or 3) who are present during outbreaks caused by a vaccine serogroup, administer 2 primary doses of MCV4-D, ideally at ages 9 months and 12 months or at least 8 weeks apart.
 - For children aged 24 months and older with 1) persistent complement component deficiency who have not been previously vaccinated; or 2) anatomic/functional asplenia, administer 2 primary doses of either MCV4 at least 8 weeks apart.
 - For children with anatomic/functional asplenia, if MCV4-D (Menactra) is used, administer at a minimum age of 2 years and at least 4 weeks after completion of all PCV doses.
 - See *MMWR* 2011;60:72–6, available at <http://www.cdc.gov/mmwr/pdf/wk/mm6003.pdf>, and Vaccines for Children Program resolution No. 6/11-1, available at <http://www.cdc.gov/vaccines/programs/vfc/downloads/resolutions/06-11mening-mcv.pdf>, and *MMWR* 2011;60:1391–2, available at <http://www.cdc.gov/mmwr/pdf/wk/mm6040.pdf>, for further guidance, including revaccination guidelines.

This schedule is approved by the Advisory Committee on Immunization Practices (<http://www.cdc.gov/vaccines/recs/acip>), the American Academy of Pediatrics (<http://www.aap.org>), and the American Academy of Family Physicians (<http://www.aafp.org>).
Department of Health and Human Services • Centers for Disease Control and Prevention



SIDS Facts

What we know about SIDS:

It occurs in all kinds of families.

It occurs in seemingly healthy infants.

It has nothing to do with race or economic status.

It occurs most often in fall and winter months.

Most deaths happen before six months of age, and especially between one and four months of age.

It is the leading cause of death in infants between one month and one year of age.

It often happens quickly during sleep, and the infant shows no signs of suffering.

It occurs most often among boys, premature and low-birth weight infants, twins, and triplets.

It is determined as the cause of death only after all other causes have been eliminated through an autopsy, a thorough investigation of the death scene, and a review of the family history.

No one knows its cause.

Safe sleep top 10:

Always place the infant on his or her back to sleep, for naps and at night; it is the safest.

Always place the infant on a firm sleep surface, such as on a safety-approved crib mattress, covered by a fitted sheet. Never place the infant to sleep on pillows, quilts, sheepskins, or other soft surfaces.

Always keep soft objects, toys, and loose bedding out of the infant's sleep area. Do not use pillows, blankets, quilts, sheepskins, and pillow-like crib bumpers in the infant's sleep area, and keep any other items away from the infant's face.

Never allow smoking around the infant.

Keep the infant's sleep area close to, but separate from, where you and others sleep. The infant should not sleep in a bed or on a couch or armchair with adults or other children, but he or she can sleep in the same room as you. If you bring the infant into bed with you to breastfeed, put him or her back in a separate sleep area, such as a bassinet, crib, cradle, or a bedside co-sleeper (infant bed that attaches to an adult bed) when finished.

Think about using a clean, dry pacifier when placing the infant down to sleep, but do not force the infant to take it. (If you are breastfeeding, wait until the infant is one month old or is used to breastfeeding before using a pacifier.)

Do not let the infant overheat during sleep. Dress the infant in light sleep clothing, and keep the room at a temperature that is comfortable for an adult.

Avoid products that claim to reduce the risk of SIDS because most have not been tested for effectiveness or safety.

Do not use home monitors to reduce the risk of SIDS. If you have questions about using monitors for other conditions, talk to your health care provider.

To reduce the chance that flat spots will develop on the infant's head from too much time on his or her back, provide "tummy time" when the infant is awake and someone is watching; change the direction that the infant lies in the crib from one week to the next; and avoid too much time in car seats, carriers, and bouncers.